

Design and Development of Fe-Catalyzed Intra- and Intermolecular Carbofunctionalization of Vinyl Cyclopropanes

Lei Liu, Wes Lee, Mingbin Yuan, Chris Acha, Michael B. Geherty, Brandon Williams, and Osvaldo Gutierrez*[‡]

[‡] Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742, United States

ABSTRACT: Design and implementation of the first (asymmetric) Fe-catalyzed intra- and intermolecular difunctionalization of vinyl cyclopropanes (VCPs) with alkyl halides and aryl Grignard reagents has been realized via a mechanistically driven approach. Mechanistic studies support the diffusion of the radicals intermediates out of the solvent cage to participate in an intra- or intermolecular radical cyclization/ring-opening followed by re-entering the Fe radical cross-coupling cycle to undergo C(sp²)-C(sp³) bond formation. Overall, we provide new design principles for Fe-mediated radical processes and underscore the potential of using combined computations and experiments to accelerate the development of challenging transformations.


INTRODUCTION

Iron-catalyzed C-C cross-coupling reactions have attracted much attention due to higher abundance, cost-effective, and lower toxicity of iron in comparison to precious transition metals.¹ Methods for ligand-supported (e.g., N-heterocyclic, bisphosphine, and diamines) and ligand-free systems for iron-catalyzed C-C cross-coupling reactions using C(sp), C(sp²), and C(sp³) partners have been developed.² In particular, bisphosphine-iron systems have emerged as highly versatile and promising manifolds for the formation of new C-C bonds with a range of organometallic nucleophiles including Mg- (Kumada),³ Zn- (Negishi),⁴ B- (Suzuki-Miyaura),⁵ and Al⁶ with alkyl halides and redox active esters.^{4a} Electron-poor vinyl cyclopropanes have also been used as π -coupling partners in Fe-catalyzed C(sp²)-C(sp³) bond formations (**Scheme 1A**). In particular, Fürstner used low valent iron ferrates to promote C-C bond formation using aryl Grignard reagents and vinyl cyclopropanes (VCP).⁷ Plietker used a low valent electronic rich ferrate complex (Bu₄N[Fe(CO)₅(NO)]) to promote coupling between a range of acidic pronucleophiles and electron-deficient VCPs.⁸ Despite these advancements, to the best of our knowledge, there are only two reports of asymmetric iron-catalyzed cross-couplings: between aryl Grignards or lithium aryl borates as nucleophiles and α -halo esters as electrophile (**Scheme 1A**).^{9,10} Thus, development of new (asymmetric) iron-catalyzed radical cascade/C(sp²)-C(sp³) cross-coupling reactions will expand the synthetic toolbox and lead to an increase in diversification of carbocycles.¹¹ Herein, we used a mechanistically-guided approach to design and develop new Fe-catalyzed *intra-* and *inter-*molecular dicarbofunctionalization of synthetically versatile vinyl cyclopropanes. (**Scheme 1B**).^{12, 13}

Contributing to the scarcity of iron-catalyzed asymmetric methods is likely due to the fact that mechanistic details of iron-catalyzed cross-coupling are not well-understood, in comparison to palladium systems.¹⁴ Pioneering mechanistic

studies by Kochi¹⁵ in the 1970s and more recent reports by Bedford,¹⁶ Nakamura,¹⁷ Norrby,¹⁸ Fürstner,¹⁹ Tonzetich,²⁰

Scheme 1. Design and development of Fe-catalyzed intra- and inter-molecular functionalization of vinyl cyclopropanes via new radical cascade reaction.




Koszinowski,²¹ and Neidig²² have led to greater understanding of these transformations. In 2017, parallel quantum mechanical studies in our lab²³ and by Morokuma²⁴ were reported on the mechanism of chiral bisphosphine cross-coupling reactions between α -chloro esters and aryl Grignard reagents. These studies revealed a mechanism involving halogen abstraction by aryl Fe(I) leading for an alkyl radical and halo aryl Fe(II) species (**Scheme 1C**; circled). In turn, these two species could combine leading to Fe(III) intermediate which will then undergo reductive elimination leading to the desired product. More experimental studies (i.e., spectroscopic and kinetic) are needed to assess the validity of the computational models and the mechanism likely depends on subtle changes to the alkyl halide, Grignard, and ligand structures. Nonetheless, based on these mechanistic studies, we envisage diverting the reactivity from the Fe radical cross-coupling catalytic cycle (black) to a programmed *intra*- and *inter*molecular radical cascade (red) with vinyl cyclopropanes leading to a new alkyl radical that could then re-enter the catalytic cycle and undergo C(sp²)-C(sp³) bond formation. Given that most of the transition-metal catalyzed cascade reactions terminate with C-H bond formation²⁵ and fail to control the stereoselectivity at the termination step, if successful, this approach could lead to rapid molecular complexity.

RESULTS AND DISCUSSION

Stereoselective *Intra*-molecular dicarbofunctionalization of VCP. Li's group demonstrated the use of cyclopropyl olefins to promote a radical alkylation, ring-opening, *intra*molecular arylation cascade reaction under photoredox conditions.²⁶ Fu and co-workers reported that chiral nickel catalysts could achieve enantioselective cross-couplings of a wide range of racemic alkyl halides (as radical precursors) with organometallic nucleophiles²⁷ including stereoconvergent radical cyclization/arylation.²⁸ We hypothesize that, upon radical formation and in the presence of a pendant vinyl cyclopropane, (**Scheme 1C**; right), we could diverge reactivity from the cross-coupling cycle to promote an *intra*-molecular radical cascade reaction. Specifically, we envisage a 5-*exo*-trig cyclization outcompeting radical rebound to aryl Fe species and *diverge* reactivity towards *Fe*-catalyzed dicarbofunctionalization of vinyl

Scheme 2. Energetics for the in-cage (blue) and out-of-cage (red) arylation computed at the UPBEPBE/6-311+G(d,p)-SDD(Fe)-THF(SMD)//UB3LYP/6-31G(d) levels of theory.




cyclopropanes. As shown in **Scheme 2**, quantum mechanical calculations support our hypothesis. Specifically, the barrier for radical 5-*exo*-cyclization (via **TS_{A-B}**) to form **B•** is 6.0 kcal/mol *lower* in energy than the barrier for radical rebound transition state **TS_{A-P1}** that will lead to cross-coupling product **P1** (7.1 kcal/mol vs. 13.1 kcal/mol, respectively). In turn, the kinetically-favored cyclic alkyl radical **B•** will then undergo radical ring-opening (barrier is only 1.6 kcal/mol) to form the

thermodynamically favored alkyl radical **C•**. Finally, **C•** could then re-enter the iron cross-coupling cycle and undergo C(sp²)-C(sp³) bond formation (via **TS_{C-P2}**; barrier is 14.8 kcal/mol from **C•**) leading, after reductive elimination (not shown), to the radical cascade product **P2**. Overall, these calculations suggest that the radical cyclization/ring-opening/cross-coupling pathway (leading to **P2**) is kinetically favored over cross-coupling pathway (leading to **P1**). Moreover, we anticipate that the chiral aryl iron species could control the final radical coupling step (with **C•**) and permit high-levels of stereocontrol.²⁹

Gratifyingly, we found that we can diverge reactivity towards the *asymmetric intra*-molecular difunctionalization of vinyl cyclopropanes using **1** as substrate (**Scheme 3**). Specifically, using the standard conditions for Fe-catalyzed α -arylation,³⁰ the reaction of **1** with aryl Grignard gave a mixture of diastereomeric radical cascade products **2a** (see Supporting Information). To determine if the chiral iron species controlled the stereochemistry of terminating C(sp²)-C(sp³) arylation (green), we subjected **2a** to a one-pot procedure of ozonolysis followed by reduction by NaBH₄ that led to the corresponding product **3a** in good yields and enantioselectivities. Notably, as expected from radical cyclization event in the absence of chiral iron species (**Scheme 2**), the corresponding *racemic* cyclopentane fragment **3b'** was isolated in 77% yield in 4:1 dr. Moreover, as shown in **Scheme 3B**, the aryl Grignard scope is broad. Both electron-poor and electron-rich aryl Grignards participate in the diverted Fe-catalyzed *intra*-molecular dicarbofunctionalization of vinyl cyclopropanes to give the desired

Scheme 3. Design and application of asymmetric Fe-catalyzed *intra*-molecular dicarbofunctionalization of vinyl cyclopropanes.




^a O₃, CH₂Cl₂/MeOH -78 °C, 5 min then NaBH₄ (10 equiv) -78 °C to rt, 1 h.

products in good yields (up to 82% over three steps) and enantioselectivities (up to 90:10 er). Highly electron-deficient (i.e., 3,4,5-trifluorophenyl **3h**) and even sterically congested (e.g., *ortho*-methyl aryl **3i** and 1-naphthyl **3k**) aryl Grignard reagents

formed the desired products in good enantioselectivities (77:23 to 90:10 er) albeit lower yields (8-50%). However, at this moment, this method is limited to 5-*exo*-trig radical cyclization. Lengthening the tether by an extra methylene group lead to formation of the Fe radical cross-coupled product and no products from diverged 6-*exo*-trig radical cyclization are observed (See Supporting Information). Presumably the much higher energy barrier to undergo radical 6-*exo*-trig cyclization in comparison to 5-*exo*-trig radical cyclization prevents formation of radical cascade/arylation (see Figure S9 in the Supporting Information for energetics).

Design and application of *Inter-molecular dicarbofunctionalization of vinyl cyclopropanes.* Given the scarcity of stereoconvergent transition metal-catalyzed radical cyclization-arylation cascades, our proof-of-principle results with *intramolecular* functionalization of VCPs (**Scheme 2** and **3**) represent an attractive strategy towards this unmet need.³¹ We envisage that by tuning the properties of alkyl halide, we could divert reactivity towards *inter-molecular* difunctionalization of VCPs. Specifically, we hypothesize that sterically hindered *tertiary* alkyl halides will lead to *higher* barriers for *Fe radical coupling* and, instead, favor *inter-molecular* radical addition to the vinyl cyclopropane (**Scheme 4A**). In turn, the incipient radical could then undergo cyclopropyl ring-opening and re-enter the Fe radical cross-coupling cycle to undergo C(sp²)-C(sp³) arylation at the least sterically hindered site (*vide infra*).

Scheme 4. Reaction Scope on Aryl Grignard of Fe-catalyzed intermolecular dicarbofunctionalization of vinyl cyclopropanes.




^a Reactions were performed on a 0.20 mmol scale. ^b ArMgBr was added dropwise via syringe pump over 4 h. ^c Isolated yield.

Previous work by Fürstner and Plietker found that Fe-catalyzed carbofunctionalization of vinyl cyclopropanes (VCPs) terminates with C-H bond formation (**Scheme 1A**).^{7, 8} Thus, if successful, this approach will expand the range of Fe-

catalyzed transformations using vinyl cyclopropanes as valuable synthons in chemical synthesis.

We initiated our optimization studies using sterically hindered *tert*-butyl 2-bromo-2-methylpropanoate **4**, (1-cyclopropylvinyl)benzene **5** and aryl 4-fluorophenylmagnesium bromide **6** were selected as model substrates (**Scheme 4**). Gratifyingly, after extensive ligand screening using both bisphosphine, monophosphine, and diamine ligands (See Table S2 in the Supporting Information) and optimization studies, we identified acyclic 1,2-bis(dimethylphosphino)ethane ligand **L11** suitable to promote the desired transformation forming compound **7a** in 82% isolated yield and 3.6:1 *E/Z* ratio (**Scheme 4B**). We also performed quantum mechanical calculations on the intermolecular dicarbofunctionalization of this system, and the calculated result is consistent with our hypothesis and the experiment result (see Figure S10 in the Supporting Information for energetics). Specifically, *tertiary* alkyl radical is favored to undergo *inter-molecular* radical addition (with barrier of 14.0 kcal/mol) to the vinyl cyclopropane over radical coupling with Fe (with barrier of 18.2 kcal/mol).

Scheme 5. Vinyl and Alkyl Halide Reaction Scope.




^a Reactions were performed on a 0.20 mmol scale. ^b ArMgBr was added dropwise via syringe pump over 4 h. ^c Isolated yield.

With optimized conditions in hand, we next investigated the reaction scope of this transformation. As shown in **Scheme 4B**, the scope of this Fe-catalyzed 3-component dicarbofunctionalization is fairly broad with respect to the Grignard nucleophile. Specifically, the reaction tolerated both electron-withdrawing and electron-donating aryl nucleophiles forming the desired 1,5-dicarbofunctionalization products in 61-92% yield. Notably, the reaction tolerated sterically hindered *ortho*-substituted aryl Grignards (**7h**, **7k**, **7l**, **7m**) that proved problematic in previous Fe-catalyzed cross-coupling reactions.³² To highlight the versatility of this method, we used both *vinyl* and *heteroaryl* nucleophiles and, gratifyingly, provided the desired products **7n** and **7o**, in good yield, 70% and 65% yield, respectively. We further explored the scope of the vinyl cyclopropane and alkyl radical precursors (**Scheme 5**). Installing electron-rich or electron-poor aryl groups in the vinyl cyclopropane moiety (R⁴) formed the desired products with good yields (**Scheme 5A**). Further, the reaction tolerates medicinally relevant heterocyclic aryl groups (**7s**, **7t**) and the substituted cyclopropanes (R⁵) were suitable for this catalytic system, affording **7u** and **7v** in 67% and 34% yield, respectively. It is

worth noting that alkyl VCPs is beyond the reach of the present Fe-based system such as **7w**. We next examined the substrate scope in regard to the alkyl radical precursor (**Scheme 5B**). Notably, we found that both alkyl fluorinated radical precursors are effective partners in the Fe-catalyzed intermolecular dicarbofunctionalization of VCPs yielding the desired products **7x** and **7y** in 42% and 75% yield, respectively. Further, we observed a much higher *E/Z* ratio for **7y** although the origin of this high selectivity is currently unknown. Finally, unactivated 3° alkyl radical precursors also formed to the desired 3-component **7z** and **7za** products. Pleasingly, as predicted by computations, preliminary results (**Scheme 6**) demonstrate that the chiral iron species could control enantioselectivity at the terminal site of C-C formation in the presence of an ester moiety although, at the current stage, this reaction suffers from low yields.

Scheme 6. Preliminary results of asymmetric Fe-catalyzed intermolecular dicarbofunctionalization of vinyl cyclopropanes.



^a Reactions were performed on a 0.20 mmol scale. ^b ArMgBr was added dropwise via syringe pump over 4 h. ^c Isolated yield. ^d The enantiomeric ratio (er) value of (*E*)-**8** was determined using chiral HPLC analysis.

CONCLUSIONS

In summary, Fe-catalyzed *intra*- and *inter*-molecular dicarbofunctionalization of vinyl cyclopropanes have now been realized. In particular, we have used a mechanistically driven (computational and experimental) approach to diverge reactivity of the alkyl radical from the Fe radical cross-coupling cycle to undergo *intra*- and *inter*-molecular radical cascade reaction with vinyl cyclopropanes. In turn, the incipient alkyl radical could then re-enter the Fe cross-coupling cycle and undergo (stereoselective) C(sp²)-C(sp³) bond formation. We anticipate the intermolecular 3-component Fe-catalyzed dicarbofunctionalization reaction to impact the synthesis medicinally relevant molecules. Ongoing work is focused on expanding this strategy to a wide range of alkyl radicals and π -coupling partners participating in diverged Fe-catalyzed radical cascade/cross-coupling reactions and using quantum mechanical calculations to optimize asymmetric variations.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental and computational details, coordinates, and spectral data (PDF)

Crystallographic data (CIF)

AUTHOR INFORMATION

Corresponding Author

* Email: ogs@umd.edu

ORCID

Lei Liu: 0000-0002-1708-9246

Wes Lee: 0000-0002-8754-6481

Osvaldo Gutierrez: 000-0001-8151-7519

Mingbin Yuan: 0000-0002-4658-4799

Chris Acha: 0000-0001-8032-2330

Michael B. Geherty: 0000-0002-0218-171X

Brandon Williams: 0000-0003-0179-7587

ACKNOWLEDGMENT

We are grateful for the financial support by the NSF (CAREER 1751568). O.G. is grateful to the University of Maryland College Park for start-up funds and computational resources from UMD Deepthought2 and MARCC/BlueCrab HPC clusters and XSEDE (CHE160082 and CHE160053). We are grateful to Prof. Lai-Xi Wang (UMD) and Prof. Donald Watson (U. Delaware) for the use of the prep HPLC and polarimeter, respectively. Finally, we were grateful to Christopher Kelly, Prof. Martinez-Solorio (Drexel), Prof. Phillip DeShong (UMD), Prof. Davis (UMD), and Prof. Andrei Vedernikov (UMD) for helpful suggestions. We also acknowledge the Analytical NMR Service & Research Center (UMD) for NMR service/help provided by Dr. YindeWang and Dr. Fu Chen. We also thank Dr. Yue Li (UMD) for help with the HRMS and Peter Y. Zavaliji for help with X-ray diffraction.

REFERENCES

- For a recent review, see: Piontek, A.; Bisz, E.; Szostak, M. Iron-Catalyzed Cross-Couplings in the Synthesis of Pharmaceuticals: In Pursuit of Sustainability. *Angew. Chem. Int. Ed.* **2018**, *57*, 11116–11128.
- For selected reviews, see: (a) Cassani, C.; Bergonzini, G.; Walentin, C.-J. Active Species and Mechanistic Pathways in Iron-Catalyzed C–C Bond-Forming Cross-Coupling Reactions. *ACS Catal.* **2016**, *6*, 1640–1648. (b) Bedford, R. B. How Low Does Iron Go? Chasing the Active Species in Fe-Catalyzed Cross-Coupling Reactions. *Acc. Chem. Res.* **2015**, *48*, 1485–1493. (c) Bauer, I.; Knölker, H.-J. Iron Catalysis in Organic Synthesis. *Chem. Rev.* **2015**, *115*, 3170–3387. (d) Czaplik, W. M.; Mayer, M.; Cvengros, J.; von Wangelin, A. J. Coming of age: sustainable iron-catalyzed cross-coupling reactions. *ChemSusChem.* **2009**, *2*, 396–417. (e) Sherry, B. D.; Fürstner, A. The Promise and Challenge of Iron-Catalyzed Cross Coupling. *Acc. Chem. Res.* **2008**, *41*, 1500–1511.
- (a) Sun, C.-L.; Krause, H.; Fürstner, A. A Practical Procedure for Iron-Catalyzed Cross-Coupling Reactions of Sterically Hindered Aryl-Grignard Reagents with Primary Alkyl Halides. *Adv. Synth. Catal.* **2014**, *356*, 1281–1291. (b) Kawamura, S.; Nakamura, M. Ligand-controlled Iron-catalyzed Cross Coupling of Benzylic Chlorides with Aryl Grignard Reagents. *Chem. Lett.* **2013**, *42*, 183–185. (c) Hatakeyama, T.; Fujiwara, Y.; Okada, Y.; Itoh, T.; Hashimoto, T.; Kawamura, S.; Ogata, K.; Takaya, H.; Nakamura, M. Kumada-Tamao-Corriu coupling of alkyl halides catalyzed by an iron-bisphosphine complex. *Chem. Lett.* **2011**, *40*, 1030–1032. (d) Hatakeyama, T.; Okada, Y.; Yoshimoto, Y.; Nakamura, M. Tuning Chemoselectivity in Iron-Catalyzed Sonogashira-Type Reactions Using a Bisphosphine Ligand with Peripheral Steric Bulk: Selective Alkynylation of Nonactivated Alkyl Halides. *Angew. Chem., Int. Ed.* **2011**, *50*, 10973–10976. (e) Dongol, K. G.; Koh, H.; Sau, M.; Chai, C. L. L. Iron-catalyzed sp³-sp³ cross-coupling reactions of unactivated alkyl halides with alkyl Grignard reagents. *Adv. Synth. Catal.* **2007**, *349*, 1015–1018.

- (a) Toriyama, F.; Cornella, J.; Wimmer, L.; Chen, T.-G.; Dixon, D. D.; Creech, G.; Baran, P. S. Redox-Active Esters in Fe-catalyzed C-C Coupling. *J. Am. Chem. Soc.* **2016**, *138*, 11132–11135. (b) Bedford, R. B.; Carter, E.; Cogswell, P. M.; Gower, N. J.; Haddow, M. F.; Harvey, J. N.; Murphy, D. M.; Neeve, E. C.; Nunn, J. Simplifying Iron-Phosphine Catalysts for Cross-Coupling

- Reactions. *Angew. Chem. Int. Ed.* **2013**, *52*, 1285–1288. (c) Adams, C. J.; Bedford, R. B.; Carter, E.; Gower, N. J.; Haddow, M. F.; Harvey, J. N.; Huwe, M.; Cartes, M. A.; Mansell, S. M.; Mendoza, C.; Murphy, D. M.; Neeve, E. C.; Nunn, J. Iron(I) in Negishi Cross-Coupling Reactions. *J. Am. Chem. Soc.* **2012**, *134*, 10333–10336. (d) Hatakeyama, T.; Kondo, Y.; Fujiwara, Y.; Takaya, H.; Ito, S.; Nakamura, E.; Nakamura, M. Iron-catalysed fluoroaromatic coupling reactions under catalytic modulation with 1,2-bis(diphenylphosphino)benzene. *Chem. Commun.* **2009**, 1216–1218. (e) Bedford, R. B.; Huwe, M.; Wilkinson, M. C. Iron-catalysed Negishi coupling of benzyl halides and phosphates. *Chem. Commun.* **2009**, 600–602.
- 5 (a) Nakagawa, N.; Hatakeyama, T.; Nakamura, M. Iron-catalyzed Suzuki–Miyaura Coupling Reaction of Unactivated Alkyl Halides with Lithium Alkynylborates. *Chem. Lett.* **2015**, *44*, 486–488. (b) Hatakeyama, T.; Hashimoto, T.; Kathiriarachchi, K. K. A. D. S.; Zenmyo, T.; Seike, H.; Nakamura, M. Iron-Catalyzed Alkyl-Alkyl Suzuki–Miyaura Coupling. *Angew. Chem. Int. Ed.* **2012**, *51*, 8834–8837. (c) Hashimoto, T.; Hatakeyama, T.; Nakamura, M. Stereospecific Cross-Coupling between Alkenylboronates and Alkyl Halides Catalyzed by Iron–Bisphosphine Complexes. *J. Org. Chem.* **2011**, *77*, 1168–1173. (d) Hatakeyama, T.; Hashimoto, T.; Kondo, Y.; Fujiwara, Y.; Seike, H.; Takaya, H.; Tamad, Y.; Ono, T.; Nakamura, M. Iron-Catalyzed Suzuki–Miyaura Coupling of Alkyl Halides. *J. Am. Chem. Soc.* **2010**, *132*, 10674–10576.
- 6 (a) Kawamura, S.; Kawabata, T.; Ishizuka, K.; Nakamura, M. Iron-catalysed cross-coupling of halohydrins with aryl aluminium reagents: a protecting-group-free strategy attaining remarkable rate enhancement and diastereoselection. *Chem. Commun.* **2012**, *48*, 9376–9378. (b) Kawamura, S.; Ishizuka, K.; Takaya, H.; Nakamura, M. The first iron-catalysed aluminium-variant Negishi coupling: critical effect of co-existing salts on the dynamic equilibrium of arylaluminium species and their reactivity. *Chem. Commun.* **2010**, *46*, 6054–6056.
- 7 Sherry, B. D.; Fürstner, A. Iron-catalyzed addition of Grignard reagents to activated vinyl cyclopropanes. *Chem. Comm.* **2009**, 7116–7118.
- 8 Dieskau, A. P.; Holzwarth, M. S.; Plietker, B. Fe-Catalyzed Allic C–C Bond Activation: Vinylcyclopropanes As Versatile α , α 3, δ 5-Synthods in Traceless Allylic Substitutions and [3+2]-Cycloadditions. *J. Am. Chem. Soc.* **2012**, *134*, 5048–5051.
- 9 (a) Iwamoto, T.; Okuzono, C.; Adak, L.; Jin, M.; Nakamura, M. Iron-Catalysed Enantioselective Suzuki–Miyaura Coupling of Racemic Alkyl Bromides. *Chem. Commun.* **2019**, *55*, 1128–1131. (b) Jin, M.; Adak, L.; Nakamura, M. Iron-Catalyzed Enantioselective Cross-Coupling Reactions of α -Chloroesters with Aryl Grignard Reagents. *J. Am. Chem. Soc.* **2015**, *137*, 7128–7134.
- 10 For a recent review, see: Pellissier, H. Recent Developments in Enantioselective Iron-Catalyzed Transformations. *Coord. Chem. Rev.* **2019**, *386*, 1–31.
- 11 For recent reviews, see: (a) Choi, J.; Fu, G. F. Transition metal-Catalyzed Alkyl-Alkyl Bond Formation: Another Dimension in Cross-Coupling. *Science* **2017**, *356*, 6334. (b) Cherney, A. H.; Kadunce, N. T.; Reisman, S. E. Enantioselective and Enantiospecific Transition-Metal-Catalyzed Cross-Coupling Reactions of Organometallic Reagents to Construct C–C Bonds. *Chem. Rev.* **2015**, *115*, 9587–9652.
- 12 For a recent review on vinyl cyclopropanes in synthesis, see: Meazza, M.; Guo, H.; Rios, R. Synthetic applications of vinyl cyclopropane opening. *Org. Biomol. Chem.* **2017**, *15*, 2479–2490.
- 13 For iron-catalyzed functionalization of VCP, see: Chen, C.; Shen, X.; Chen, J.; Hong, X.; Lu, Z. Iron-Catalyzed Hydroboration of Vinylcyclopropanes. *Org. Lett.* **2017**, *19*, 5422–5425.
14. (a) Neidig, M. L.; Carpenter, S. H.; Curran, D. J.; DeMuth, J. C.; Fleischaer, V. E.; Iannuzzi, T. E.; Neate, P. G. N.; Sears, J. D.; Wolford, N. J. Development and Evolution of Mechanistic Understanding in Iron-Catalyzed Cross-Coupling. *Acc. Chem. Res.* **2019**, *52*, 140–150. (b) Sears, J. D.; Neate, P. G. N.; Neidig, M. L. Intermediates and Mechanism in Iron-Catalyzed Cross-Coupling. *J. Am. Chem. Soc.* **2018**, *140*, 11872–11883. (c) Liu, L.; Lee, W.; Yuan, M.; Gutierrez, O. Mechanisms of Bisphosphine Iron-Catalyzed C(sp²)-C(sp³) Cross-Coupling Reactions: Inner-Sphere or Outer-Sphere Arylation? *Comment. Inorg. Chem.* **2018**, *38*, 210–237. (d) Mako, T. L.; Byers, J. A. Recent Advances in Iron-Catalysed Cross Coupling Reactions and Their Mechanistic Underpinning. *Inorg. Chem. Front.* **2016**, *3*, 766–790.
- 15 Smith, R. S.; Kochi, J. K. Mechanistic Studies of Iron Catalysis in the Cross Coupling of Alkenyl Halides and Grignard Reagents. *J. Org. Chem.* **1976**, *41*, 502–509.
- 16 Bedford, R. B.; Brenner, P. B.; Carter, E.; Cogswell, P. M.; Haddow, M. F.; Harvey, J. N.; Murphy, D. M.; Nunn, J.; Woodall, C. H. TMEDA in Iron-Catalyzed Kumada Coupling: Amine Adduct versus Homoleptic “ate” Complex Formation. *Angew. Chem. Int. Ed.* **2014**, *53*, 1804–1808.
- 17 (a) Takaya, H.; Nakajima, S.; Nakagawa, N.; Isozaki, K.; Iwamoto, T.; Imayoshi, R.; Gower, N. J.; Adak, L.; Hatakeyama, T.; Honma, T.; Takagaki, M.; Sunada, Y.; Nagashima, H.; Hashizume, D.; Takahashi, O.; Nakamura, M. Investigation of Organoiron Catalysis in Kumada–Tamao–Corriu-Type Cross-Coupling Reaction Assisted by Solution-Phase X-ray Absorption Spectroscopy. *Bull. Chem. Soc. Jpn.* **2015**, *88*, 410–418. (b) Noda, D.; Sunada, Y.; Hatakeyama, T.; Nakamura, M.; Nagashima, H. Effect of TMEDA on Iron-Catalyzed Coupling Reactions of ArMgX with Alkyl Halides. *J. Am. Chem. Soc.* **2009**, *131*, 6078–6079.
- 18 (a) Bekhradnia, A.; Norrby, P.-O. New Insights into the Mechanism of Iron-Catalyzed Cross-Coupling Reactions. *Dalton Trans.* **2015**, *44*, 3959–3962. (b) Hedström, A.; Izakian, Z.; Vreto, I.; Wallentin, C.-J.; Norrby, P.-O. On the Radical Nature of Iron-Catalyzed Cross-Coupling Reactions. *Chem. - Eur. J.* **2015**, *21*, 5946–5953. (c) Kleimark, J.; Hedström, A.; Larsson, P.-F.; Johansson, C.; Norrby, P.-O. Mechanistic Investigation of Iron-Catalyzed Coupling Reactions. *ChemCatChem* **2009**, *1*, 152–161. (d) Hedström, A.; Bollmann, U.; Bravidor, J.; Norrby, P.-O. Iron-Catalyzed Coupling of Aryl Grignard Reagents with Alkyl Halides: A Competitive Hammett Study. *Chem. - Eur. J.* **2011**, *17*, 11991–11993.

- 19 (a) Casitas, A.; Rees, J. A.; Goddard, R.; Bill, E.; DeBeer, S.; Fürstner, A. Two Exceptional Homoleptic Iron(IV) Tetraalkyl Complexes. *Angew. Chem. Int. Ed.* **2017**, *56*, 10108–10113. (b) Fürstner, A.; Krause, H.; Lehmann, C. W. Unusual Structure and Reactivity of a Homoleptic “Super-Ate” Complex of Iron: Implications for Grignard Additions, Cross-Coupling Reactions, and Kharasch Deconjugation. *Angew. Chem. Int. Ed.* **2006**, *45*, 440–444.
- 20 Przyojski, J. A.; Veggeberg, K. P.; Arman, H. D.; Tonzetich, Z. J. Mechanistic Studies of Catalytic Carbon–Carbon Cross-Coupling by Well-Defined Iron NHC Complexes. *ACS Catal.* **2015**, *5*, 5938–5946.
- 21 (a) Parchomyk, T.; Demeshko, S.; Meyer, F.; Koszinowski, K. Oxidation States, Stability, and Reactivity of Organoferrate Complexes. *J. Am. Chem. Soc.* **2018**, *140*, 9709–9720. (b) Parchomyk, T.; Koszinowski, K. Ate complexes in iron-catalyzed cross-coupling reactions. *Chem. - Eur. J.* **2016**, *22*, 15609–15613.
- 22 (a) Carpenter, S. H.; Baker, T. M.; Muñoz III, S. B.; Brennessel, W. W.; Neidig, M. L. Multinuclear iron–phenyl species in reactions of simple iron salts with PhMgBr: identification of Fe₄(μ-Ph)₆(THF)₄ as a key reactive species for cross-coupling catalysis. *Chem. Sci.* **2018**, *9*, 7931–7939. (b) Muñoz III, S. B.; Daifuku, S. L.; Sears, J. D.; Baker, T. M.; Carpenter, S. H.; Brennessel, W. W.; Neidig, M. L. The N-Methylpyrrolidone (NMP) Effect in Iron-Catalyzed Cross-Coupling with Simple Ferric Salts and MeMgBr. *Angew. Chem. Int. Ed.* **2018**, *57*, 6496–6500. (c) Fleischauer, V. E.; Muñoz III, S. B.; Neate, P. G. N.; Brennessel, W. W.; Neidig, M. L. NHC and nucleophile chelation effects on reactive iron(II) species in alkyl–alkyl cross-coupling. *Chem. Sci.* **2018**, *9*, 1878–1891. (d) Kneebone, J. L.; Brennessel, W. B.; Neidig, M. L. Intermediates and Reactivity in Iron-Catalyzed Cross-Couplings of Alkynyl Grignards with Alkyl Halides. *J. Am. Chem. Soc.* **2017**, *139*, 6988–7003. (e) Muñoz, S. B., III; Daifuku, S. L.; Brennessel, W. W.; Neidig, M. L. Isolation, Characterization, and Reactivity of Fe₈Me₁₂: Kochi's S = 1/2 Species in Iron-Catalyzed Cross-Couplings with MeMgBr and Ferric Salts. *J. Am. Chem. Soc.* **2016**, *138*, 7492–7495. (f) Daifuku, S. L.; Kneebone, J. L.; Snyder, B. E. R.; Neidig, M. L. Iron(II) Active Species in Iron-Bisphosphine Catalyzed Kumada and Suzuki-Miyaura Cross-Couplings of Phenyl Nucleophiles and Secondary Alkyl Halides. *J. Am. Chem. Soc.* **2015**, *137*, 11432–11444. (g) Al-Afyouni, M. H.; Fillman, K. L.; Brennessel, W. W.; Neidig, M. L. Isolation and Characterization of a Tetramethyliron(III) Ferrate: An Intermediate in the Reduction Pathway of Ferric Salts with MeMgBr. *J. Am. Chem. Soc.* **2014**, *136*, 15457–15460. (h) Daifuku, S. L.; Al-Afyouni, M. H.; Snyder, B. E. R.; Kneebone, J. L.; Neidig, M. L. A Combined Mossbauer, Magnetic Circular Dichroism, and Density Functional Theory Approach for Iron Cross-Coupling Catalysis: Electronic Structure, In Situ Formation, and Reactivity of Iron-Mesityl-Bisphosphines. *J. Am. Chem. Soc.* **2014**, *136*, 9132–9143.
- 23 Lee, W.; Zhou, J.; Gutierrez, O. Mechanism of Nakamura's Bisphosphine Iron-Catalyzed Asymmetric C(sp²)-C(sp³) Cross-Coupling Reaction: The Role of Spin in Controlling Arylation Pathway. *J. Am. Chem. Soc.* **2017**, *139*, 16126–16133.
- 24 Sharma, A. K.; Sameera, W. M. C.; Jin, M.; Adak, L.; Okuzono, C.; Iwamoto, T.; Kato, M.; Nakamura, M.; Morokuma, K. DFT and AFIR Study on the Mechanism and the Origin of Enantioselectivity in Iron-Catalyzed Cross-Coupling Reactions. *J. Am. Chem. Soc.* **2017**, *139*, 16117–16125.
- 25 For selected reviews, see: (a) Plesniak, M. P.; Huang, H.-M.; Procter, D. J. Radical cascade reactions triggered by single electron transfer. *Nat. Rev. Chem.* **2017**, *1*, 0077. (b) Quiclet-Sire, B.; Zard, S. Z. Some aspects of radical cascade and relay reactions. *Proc. R. Soc. A* **2017**, *473*, 20160859. (c) For a recent example of radical cascade in total synthesis, see: Brill, Z. G.; Grover, H. K.; Maimone, T. J. Enantioselective synthesis of an ophiobolin sesterterpene via a programmed radical cascade. *Science* **2016**, *352*, 1078–1082. (d) Wille, U. Radical Cascades Initiated by Intermolecular Radical Addition to Alkynes and Related Triple Bond Systems. *Chem. Rev.* **2013**, *113*, 813–853. (e) Nicolaou, K. C.; Edmonds, D. J.; Bulger, P. G. Cascade Reactions in Total Synthesis. *Angew. Chem. Int. Ed.* **2006**, *45*, 7134–7186.
- 26 Li, J.; Chen, J.; Jiao, W.; Wang, G.; Li, Y.; Cheng, X.; Li, G. Difluoroalkylation/C–H Annulation Cascade Reaction Induced by Visible-Light Photoredox Catalysis. *J. Org. Chem.* **2016**, *81*, 9992–10001.
- 27 Fu, G. C. Transition-Metal Catalysis of Nucleophilic Substitution Reactions: A Radical Alternative to S_N1 and S_N2 Processes. *ACS Cent. Sci.* **2017**, *3*, 692–700.
- 28 Cong, H.; Fu, G. C. Catalytic Enantioselective Cyclization/Cross-Coupling with Alkyl Electrophiles. *J. Am. Chem. Soc.* **2014**, *136*, 3788–3791.
- 29 For computational details, see Supporting Information.
- 30 Liu, L.; Lee, W.; Zhou, J.; Bandyopadhyay, S.; Gutierrez, O. Radical-clock α-halo-esters as mechanistic probes for bisphosphine iron-catalyzed cross-coupling reactions. *Tetrahedron* **2019**, *75*, 129–136.
- 31 Potential challenges include: a) diffusion-control rates of radical/radical recombination, b) H-atom abstraction from solvent, non-stereoselective arylation with ferric species, c) single electron transfer (SET) leading to carbocationic (or anionic) pathways, and, inter alia, d) arylation in the outer-sphere of the chiral iron species that prevents high-levels of stereocontrol in the fragment-coupling step. Further, the presumed intercepted alkyl Fe(III) intermediate could also undergo.
- 32 Jin, M.; Nakamura, M. Iron-catalyzed Chemoselective Cross-coupling of α-Bromocarboxylic Acid Derivatives with Aryl Grignard Reagents. *Chem. Lett.* **2011**, *40*, 1012–1014.

